Grass and Forage Science (1986) 41, 385–387.

Author index

ARMSTRONG R.H., COMMON T.J. and SMITH H.K. The voluntary intake and *in vivo* digestibility of herbage harvested from indigenous hill plant communities, 53.

BAKER A-M.C. and LEAVER J.D. Effect of stocking rate in early season on dairy cow performance and sward characteristics, 333.

BAKER A-M.C. and YOUNGER A. The effect of temperature on the spring growth of perennial ryegrass at three contrasting sites (Research Note), 175.

BARTHRAM G.J. *see* KING J.

BARTIAUX-THILL N. and OGER R. The indirect estimation of the digestibility in cattle of herbage from Belgian permanent pasture, 269.

BOLTON R. *see* RANGELY A.

BOYD A.G. *see* FRAME J.

BRAY R.A. *see* MINSON D.J.

CHAMBERLAIN D.G., THOMAS P.C. and QUIG J. Utilization of silage nitrogen in sheep and cows: amino acid composition of duodenal digesta and rumen microbes, 31.

CLEMENTS R.O. *see* MOORE D.

COLLISON M. *see* GOODMAN P.J.

COMMON T.G. *see* ARMSTRONG R.H.

DOUGLAS J.A. The production and utilization of lucerne in New Zealand (Review Paper), 81.

DOVE H. and MCCORMACK H.A. Estimation of the ruminal degradation of forage rape after incubation in nylon bags in the rumen of sheep, 129.

DOYLE, C.J., RIDOUT M.S., MORRISON J. and EDWARDS C. Predicting the response of perennial ryegrass to fertilizer nitrogen in relation to cutting interval, climate and soil, 303.

EDWARDS C. *see* DOYLE C.J.

ELLIS DAVIES W. *see* EVANS D.R.

EVANS D.R., WILLIAMS T.A. and ELLIS DAVIES W. Potential seed yield of white clover varieties, 221.

EVANS D.R., THOMAS T.A., WILLIAMS T.A. and ELLIS DAVIES W. Effect of fertilizers on the yield and chemical composition of pure sown white clover and on soil nutrient status, 295.

FENLON D.R. Growth of naturally occurring *Listeria* spp. in silage: a comparative study of laboratory and farm ensiled grass, 375.

FRAME J. and BOYD A.G. Effect of cultivar and seed rate of perennial ryegrass and stragic fertilizer nitrogen on the productivity of grass/white clover swards, 359.

FRASER T.J. *see* L'HULLIER P.J.

GIBB M.J. and RIDOUT M.S. The fitting of frequency distributions to height measurements on grazed swards, 247.

GOODMAN P.J. and COLLISON M. Effect of three clover varieties on growth, ^{15}N uptake and fixation by ryegrass/white clover mixtures at three sites in Wales, 191.

GORDON, F.J. The effect of system of silage harvesting and feeding on milk production.

GORDON F.J. *see* MAYNE C.S.

GORDON F J. and UNSWORTH E.F. The effects of silage harvesting system and supplementation of silage-based diets by protein and methionine hydroxy analogue on the performance of lactating cows, 1.

HAY R.K.M. and PEDERSEN K. Influence of long photoperiods on the growth of timothy (*Phleum pratense* L.) varieties from different latitudes in northern Europe, 311.

HAYES M.J. *see* WILLIAMS E.D.

HAYES M.J. and WILLIAMS E.D. Some effects of fluid-sowing, pre-germination, irrigation and soil covering on the establishment and growth of seedlings of white clover slot-seeded into permanent pasture, 151.

HENDERSON A.R. *see* SEALE D.R.

HOOPER G.E. *see* PENNING P.D.

HOPKINS A. Botanical composition of permanent grassland in England and Wales in relation to soil, environment and

management factors, 237.
L'HULLIER P.J., POPPI D.P. and FRASER T.J. Influence of structure and composition of ryegrass and prairie grass–white clover swards on the grazed horizon and diet harvested by sheep, 259.
JESS S. *see* MOWAT D.J.
JONES E.L. and ROBERTS J.E. Comparison of management techniques for evaluating Italian and hybrid ryegrass cultivars, 9.
KEER J.I., MCCLAREN R.G. and SWIFT R.G. The sulphur status of intensive grassland sites in southern Scotland. 183.
KING J., SIM E.M. and BARTHRAM G.T. A comparison of spectral reflectance and sward surface height measurements to estimate herbage mass and leaf area index in continuously stocked ryegrass pastures, 251.
LAVENDER R.H. *see* SHELDRICK R.D.
LEAVER J.D. *see* BAKER A-M.C.
LEAVER J.D. *see* ROBERTS D.J.
LIACOS L.G. and PAPANASTASIS V.P. Organic and inorganic fertilization compared in a natural Mediterranean grassland, 353.
LINZELL B.S. and MADGE D.S. Effects of pesticides and fertilizer on invertebrate populations of grass and wheat plots in Kent in relation to productivity and yield, 159.
LOWE J.F. *see* SEALE D.R.
MCADAM J.H. Growth and production of a *Cortaderia pilosa* (Graminae) community in the Falkland Islands, 281.
MCLAREN R.G. *see* KEER J.I.
MCCORMACK H.A. *see* DOVE H.
MADGE D.S. *see* LINZELL B.S.
MAYNE C.S. and GORDON F.J. The effect of harvesting system on nutrient losses during silage making. 1. Field losses, 17.
MAYNE C.S. and GORDON F.J. Effect of harvesting system on nutrient losses during silage making. 2. In-silo losses, 341.
MEIJS J.A.C. Concentrate supplementation of grazing dairy cows. 2. Effect of concentrate composition on herbage intake and milk production, 229.
MINSON D.J. and BRAY R.A. Voluntary intake and *in vivo* digestibility by sheep of five lines of *Cenchus ciliaris* selected on the basis of preference rating, 47.
MOORE D. and CLEMENTS R.O. Study of the population dynamics of dipterous stem borer larvae of ryegrass swards under rotationally grazed or cut conditons, 289.
MORRISON J. *see* DOYLE C.J.
MOWAT D.J. and JESS S. The control of leatherjackets *Tipula paludosa* Meig., in grassland by early application of insecticide, 27.
OGER R. *see* BARTIAUX-THILL N.
PAPANASTASIS V.P. *see* LIACOS L.G.
PEDERSEN K. *see* HAY R.K.M.
PENNING P.D., HOOPER G.E. and TREACHER T.T. The effect of herbage allowance on intake and performance of ewes suckling twin lambs, 199.
PETTERSON K.O. *see* SEALE D.R.
POPPI D.P. *see* L'HULLIER P.J.
QUIG J. *see* CHAMBERLAIN D.G.
RANGELY A. and BOLTON R. Lime and major nutrient fertilizers required to establish a perennial ryegrass/white clover pasture on a non-calcareous gley in the Scottish uplands, 323.
RIDOUT M.S. *see* DOYLE C.J.
RIDOUT M.S. *see* GIBB M.J.
ROBERTS H.A. Persistence of seeds of some grass species in cultivated soil, 273.
ROBERTS J.E. *see* JONES E.L.
ROBERTS D.J. and LEAVER J.D. Supplementary feeding of forage to grazing dairy cows. 3. The effect of three daytime stocking rates on the performance of cows offered grass silage overnight, 79.
SCOTT D. Coefficients for the dry-weight rank method of botanical analysis of pasture, 319.
SEALE D.R., HENDERSON A.R., PETTERSON K.O. and LOWE J.F. The effect of addition of sugar and inoculation with two commercial inoculants on the fermentation of lucerne silage in laboratory silos, 61.
SHELDRICK R.D., LAVENDER R.H. and TEWSON V.J. The effects of frequency of defoliation, date of first cut and heading date of a perennial ryegrass companion on the yield, quality and persistence of diploid and tetraploid broad red clover.
SIM E.M. *see* KING J.
SMITH H.K. *see* ARMSTRONG R.H.
STEEN R.W.J. An evaluation of effluent from grass silage as a feed for beef cattle offered silage-based diets, 39.
SWIFT R.S. *see* KEER J.I.
TEWSON V.J. *see* SHELDRICK R.D.
THOMAS P.C. *see* CHAMBERLAIN D.G.

THOMAS T.A. *see* EVANS D.R.
TREACHER T.T. *see* PENNING P.D.
UNSWORTH E.F. *see* GORDON F.J.
WILLIAMS E.D. Further studies on the effect of phosphate fertilizer and propyzamide on the growth of white clover slot-seeded into permanent pasture (Research Note), 179.
WILLIAMS E.D. *see* HAYES M.J.
WILLIAMS E.D. and HAYES M.J. Growth of white clover in eight grass species and the effects of propyzamide, a grass-suppressing herbicide, 367.
WILLIAMS T.A. *see* EVANS D.R.
YOUNGER A. *see* BAKER A-M.C.